

Medicinal Plants with Hypoglycemic and Diuretic Properties: Reducing Diuretic-Induced Hyperglycemia

Ayush Kumar^{1*}, Shobha Kumari²

¹Department of Pharmacy Practice, MM College of Pharmacy, Maharishi Markandeshwar (Deemed to be university), Mullana, Ambala

²Department of Nursing, Gautam Institute of Nursing and Para medics, Nalanda, Bihar, India

***Corresponding Author** – Ayush Kumar,

Date of Submission: 20-12-2024

Date of acceptance: 02-01-2025

ABSTRACT: Diuretic therapy, while essential for managing conditions such as hypertension and edema, often induces hyperglycaemic as a side effect, posing significant challenges for patients with diabetes or prediabetes. This review explores the potential of medicinal plants possessing diuretic and hypoglycaemic properties as a promising strategy to mitigate diuretic-induced hyperglycemia. A comprehensive analysis of relevant literature identifies several plant species known for their diuretic effects, such as *Samanea Saman*, *H. antidysentrica*, *Achyranthes Aspera*, *Taraxacum Officinale*, and *Urtica Dioica*, and their concurrent hypoglycaemic activities. These plants control glucose metabolism while preserving renal function through a variety of mechanisms that are revealed by scientific investigations into their pharmacological activities. Moreover, scientific and clinical research demonstrates the safety and effectiveness characteristics of these plants, emphasizing their potential as adjuvant medicines in medical scenarios. This synthesis aims to provide a foundation for future research and therapeutic interventions aimed at optimizing diuretic therapy while managing hyperglycemic complications effectively.

Keywords: -Diuretic, Hyperglycaemia, Hypoglycaemia

I. INTRODUCTION

Any drug that raises urine flow and thus increases the excretion of water is called a diuretic. Among the most widely prescribed medications are diuretics, most of which work by decreasing the reabsorption of sodium chloride at various nephron sites, which raises urine salt levels and therefore causes water loss^[1]. Diuretics are commonly used to treat conditions including edema, hypertension, congestive heart failure, and disorders of the kidney and

liver.^[2] Diuretics are part of therapeutic strategies to help balance fluid, help shift fluid out of the interstitium, and control fluid overload characterized as ankle swelling, ascites, and pulmonary edema. This results in significant symptom relief and improves patients' quality of life overall.^[3] Nowadays many commercial diuretic drugs are available in the market, such as the high-ceiling loop and thiazides diuretic. These commercial diuretic drugs have a variety of side effects such as hypokalemia, metabolic alkalosis, and dehydration (hypovolemia), leading to hypotension, fever, cough, unusual bleeding, hyperglycemia, excessive weight loss, nausea, and vomiting.^[2]

Drug-induced hyperglycemia is one of the global issues, increasing the risk of infections, metabolic coma, microvascular and macrovascular complications, and even death. People with certain risk factors are particularly susceptible to developing diabetes mellitus, including those with a sedentary lifestyle, a body mass index (BMI) over

27 kg/m², impaired glucose tolerance or fasting glucose, a family history of diabetes, a history of vascular disease, gestational diabetes, or at least one risk factor for metabolic syndrome. These individuals are especially vulnerable to drug-induced hyperglycemia because some medications can worsen pre-existing insulin resistance or pancreatic dysfunction.^[4]

Diuretics are widely prescribed for various cardiovascular conditions but are associated with metabolic side effects, including an increased risk of developing diabetes mellitus.^[5] The search for alternative therapies has led to exploring medicinal plants known for their diuretic actions, which may also offer hypoglycemic benefits, thus presenting a promising avenue for managing fluid retention without compromising glucose homeostasis.^[1]

II. MATERIALS AND METHODS

The current study utilized a comprehensive and systematic literature search. Various databases such as PubMed and Google Scholar were searched using terms such as diuretic, hyperglycemia, hypoglycemia, traditional uses, and medicinal plants. The identified medicinal plants were grouped based on the diuretic and hypoglycaemic actions and data from both in vitro and/or in vivo studies were compiled. This information, along with basic details about medicinal plants, was collected and carefully analyzed. Scientific names and synonyms were verified through www.plantlist.org and www.tropicos.org. References of selected articles were manually searched for additional relevant studies.

Mechanisms of diuretic-induced hyperglycemia

Thiazide diuretics are among the most commonly used antihypertensives.^[6] In a recent meta-analysis of antihypertensive trials, thiazides were associated with a higher risk of diabetes than placebo.^[7] The mechanisms behind diuretic-induced hyperglycemia appear to involve decreased insulin

production, secondary to diuretic-induced hypokalaemia.^[8,9]

Other pathways that could lead to thiazide-induced hyperglycemia include increased levels of free fatty acids, which are known to reduce insulin secretion in response to glucose- marked decreases in insulin sensitivity, and increased hepatic glucose production and/or catecholamine secretion and action ^{[10],[11]}.

Medicinal plants with dual-action

As alternate sources of treatment regimens, medicinal plants employed in traditional medicine are gaining popularity, particularly when supported by scientific proof of their clinical value. This is because several natural substances used in folk medicine are efficient, occasionally have comparatively less negative clinical effects, and are frequently reasonably priced.^[12] However, research in ethnobotany and ethnopharmacology has shown that numerous medicinal plants from various regions possess both diuretic and hypoglycaemic effects. This suggests promising avenues for discovering therapeutic agents that could effectively manage both diabetes and diuresis simultaneously.^[13]

Table No: 1.1 Plants with Both Diuretic and Hypoglycemic Activity

PLANT	FAMILY	PLANT PART WITH DIURETIC EFFECT	PLANT PART WITH HYPOGLYCAEMIC EFFECT	REF
Abutilon indicum	Malvaceae	leaf	leaf	[14,27]
Allium sativum	Alliaceae	Bulb	Bulb	[13]
Benincasa hispida	Cucurbitaceae.	fruit	fruit	[15]
Boerhavia diffusa Linn	Nyctaginaceae.	leaf	leaf	[16,28]
Centella asiatica	Apiaceae	leaf	leaf	[17]
Ipomoea aquatica	Convolvulaceae	whole plant	leaf	[18]
Xanthium strumarium	Compositae	whole plant	whole plant	[19,20]
Samanea Saman (Jacq) Merr	legumes	bark	leaf	[19,21]
Holarrhena Antidysenterica	Apocyanaceae	seed	seed	[19,22]
Achyranthes aspera linn	Amaranthaceae	whole plant	whole plant	[19,23]
Taraxacum officinale	daisy	whole plant	whole plant	[24]
Urtica dioica	Urticaceae	whole plant	whole plant	[25,26]

● **ABUTILON INDICUM**

Abutilon indicum, a member of the Malvaceae family, is commonly known as Country Mallow in English, Kanghi in Hindi, and Atibala in Sanskrit. This plant exhibits both diuretic and hypoglycemic properties.

The leaf extract of *A. indicum* was evaluated for its diuretic effect, where the aqueous extract at 400 mg/kg showed a statistically significant effect compared to the reference standard, Furosemide. Additionally, the hypoglycemic activity of *A. indicum* leaf extracts in rats was assessed, with both alcoholic and aqueous extracts demonstrating a significant reduction in blood glucose levels.^[14,27]



FigureNo:1.1 *Abutilon indicum*

● **ALLIUM SATIVUM**

Allium sativum (Meaning pungent) belongs to the Alliaceae family and genus *Allium*, and is generally known in the developing world for its characteristic flavor, as a medicinal plant, and a source of vegetable oil. The herb has hypoglycemic and diuretic properties.

Several animal studies support the effectiveness of garlic in reducing blood glucose levels in streptozotocin-induced and alloxan-induced diabetes mellitus in rats and mice. Most studies showed that garlic can lower blood glucose levels in diabetic mice, rats, and rabbits. Additionally, garlic has been reported to act as a diuretic, helping to eliminate excess body fluids, making it a potentially useful resource for conditions such as rheumatism, gout, arthritis, dropsy, and edema.^[13]



FigureNo: 1.2 *Allium sativum*

● **BENINCASA HISPIDA**

Benincasa hispida also known as Winter melon, Ash gourd, Wax gourd, White pumpkin. Belongs to the family Cucurbitaceae.

When rats were administered 0.75% v/v ethylene glycol in their drinking water to induce chronic hyperoxaluria, the concurrent oral administration of *Benincasa hispida* extract at doses of 250 and 500 mg/kg body weight for 35 days significantly reduced the urinary excretion and kidney retention levels of oxalate, protein, and calcium. Additionally, elevated serum levels of sodium, creatinine, calcium, and phosphorus were significantly reduced by the extracts. Furthermore, the stem chloroform extract of *Benincasa hispida* demonstrated significant hypoglycemic activity in normal male Wistar rats, with the maximum reduction in blood glucose levels observed at a dose of 200 mg/kg body weight.^[15]



Figure No:1.3 *Benincasa hispida*

● **BOERHAVIA DIFFUSA LINN**

Boerhavia diffusa Linn. Also known as Punarnava, Red Spiderling, or Spreading Hogweed.

Belongs to Family: Nyctaginaceae.

The chloroform extract of *Boerhavia diffusa* (BD) leaves has shown dose-dependent hypoglycemia in experimentally diabetic rats. Glibenclamide (25

µg/kg) and BD leaf extract (200 mg/kg) resulted in glucose reductions of 59.01% and 38.63%, respectively, by the fourth week. Additionally, the authors found significant diuretic activity in the BD leaf extract (a water-insoluble portion of the alcoholic extract) collected during the rainy season. Rats treated with the leaf extract (300 mg/kg) exhibited a 90.3% increase in urine volume, while the extract of leaves and flowers showed a 67.22% increase in urine volume.[16,28]



Figure No: 1.4 Boerhavia diffusa Linn

● CENTELLA ASIATICA

Centella Asiatica is also known as Gotu Kola, Indian Pennywort, or Asiatic Pennywort. Belongs to the family Apiaceae (Umbelliferae). It is a small, herbaceous, perennial plant characterized by its fan-shaped, green leaves with serrated edges. It typically grows in moist, tropical, and subtropical areas. The aqueous extracts of Centella Asiatica leaves possess good diuretic activity. The anti-diabetic properties of the leaf extract were evaluated in alloxan-induced diabetic rats by administering the extract at concentrations of 250, 500, and 1000 mg/kg. Three hours after ingestion, a reduction in blood glucose levels was observed by 32.6%, 38.8%, and 29.9%, respectively.[17]



Figure No: 1.5 Centella asiatica

● IPOMOEA AQUATICA

Ipomoea aquatica is also known as Water Spinach, Kangkong, Water Convolvulus, and Chinese Watercress. Belongs to the family Convolvulaceae. The methanol extract of Ipomoea aquatica

demonstrated notable diuretic activity in Swiss albino mice, surpassing the effects of the standard diuretic furosemide by increasing electrolyte excretion and urine volume. In addition, methanol extracts from its leaves exhibited potent hypoglycemic effects in Swiss albino mice, with doses of 200 mg/kg and 400 mg/kg significantly lowering blood glucose levels, particularly effective at 400 mg/kg. Furthermore, a boiled whole extract of Ipomoea aquatica showed oral hypoglycemic effects in healthy male Wistar rats, leading to significant reductions in serum glucose concentrations with both single (33%, $P < 0.0027$) and multiple (25%, $P < 0.02$) doses. The optimal dose was found to be 3.4 g/kg, with maximum activity observed two hours after administration.[18]



Figure No: 1.6 Ipomoea aquatic

● XANTHIUM STRUMARIUM

Xanthium strumarium, a prevalent weed in India from the Compositae family, is utilized medicinally, particularly its roots and fruits. In a study, Xanthium strumarium L. was evaluated for its diuretic activity in albino rats. The extract was administered at doses of 250 mg/kg and 500 mg/kg body weight, while furosemide, a standard diuretic, was given at 5 mg/kg body weight. The petroleum ether extract showed significant increases in diuresis, natriuresis (sodium excretion), kaliuresis (potassium excretion), and glomerular filtration rate compared to normal saline. Additionally, caffeic acid isolated from X. strumarium was studied for its antidiabetic effects in streptozotocin-induced and insulin-resistant rat models. Results indicated that caffeic acid administered intravenously at doses ranging from 0.5 to 3.0 mg/kg could lower plasma glucose levels by enhancing glucose utilization.[19,20]



Figure No:1.7 Xanthium strumarium

● SAMANEA SAMAN (JACQ) MERR

Samanea Saman (Jacq) Merr is a large, umbrella-shaped tree that can grow over 20 meters tall. It has a thick trunk approximately 1.5 meters in diameter and a wide, spreading canopy that offers ample shade. The bark is rough and deeply furrowed.

B. Komarapalayam et al. evaluated the diuretic activity of Samanea saman (Jacq) Merr bark in albino rats using the in-vivo Lipschitz test model. The study found that methanol extract of the bark at concentrations of 200 mg/kg and 400 mg/kg body weight significantly increased urine volume and electrolyte excretion compared to the control group. Furosemide, used as the standard at a dose of 20 mg/kg body weight, showed that the methanol extract of Samanea saman at higher doses might have comparable diuretic activity to furosemide. Additionally, the methanol leaf extract of Samanea saman demonstrated anti-diabetic activity in both in vitro and in vivo studies. The α -amylase inhibition showed increasing concentrations at 50 μ g/ml, 100 μ g/ml, 150 μ g/ml, 200 μ g/ml, and 250 μ g/ml. When administered to diabetic-induced rats at doses of 250 mg/kg and 500 mg/kg body weight, the extract significantly reduced blood glucose levels ($p < 0.001$) at 1 and 2 hours and ($p < 0.05$) at 30 minutes compared to the control group.^[19,21]



Figure No:1.8 Samanea Saman (Jacq) Merr

● HOLARRHENA ANTIDYSENTERICA

Holarrhena Antidysenterica belonging to the family Apocyanaceae, commonly known as bitter oleander and locally as Kurchi, is a small deciduous tree found in Himalaya and sub-Himalaya tract.

The crude extract of H. antidysenterica seeds and its fractions (n-hexane, n-butanol, and aqueous) were tested for diuretic effects in Wistar rats, using hydrochlorothiazide at 10 mg/kg body weight as the standard. The crude

aqueous ethanolic extract showed a dose-dependent increase in urine output at 30 and 100 mg/kg, indicating a diuretic effect. It also increased urine Na^+ and K^+ levels, urine volume, pH, and electrolyte levels. Additionally, a recent study reported significant recovery in diabetic rats administered 300 mg/kg and 600 mg/kg doses of the ethanolic extract of the seeds. Weekly treatments significantly decreased blood glucose, serum cholesterol, triglycerides, AST, ALT, alkaline phosphatase, urea, creatinine, and uric acid levels, while the rats' weight increased substantially.^[19,22]



Figure No:1.8 Holarrhena Antidysenterica

● ACHYRANTHES ASPERA LINN

Achyranthes aspera Linn commonly known as Amamarga in Ayurveda is a weed, in Hindi as Latjeera; belongs to the family Amaranthaceae, is an erect or procumbent, annual or perennial herb, found on roadsides, fields boundaries, and waste places as a weed.

The diuretic activity of Achyranthes aspera was evaluated using a methanolic extract of the whole plant, with the Lipschitz method and furosemide (100 mg/kg b.w.) as the standard drug. At a dose of 400 mg/kg b.w., the extract showed significant diuretic effects in rats compared to the control group. Additionally, another study reported that oral administration of 2-4 g/kg of whole plant powder produced a significant dose-related hypoglycemic effect in both normal and alloxan-treated diabetic rabbits. Both the aqueous and methanol extracts of the plant also decreased blood glucose levels in these rabbits.^[19,23]



Figure No:1.9 Achyranthes aspera Linn

● TARAXACUM OFFICINALE

The dandelion, or *Taraxacum officinale* (G.H. Weber ex Wiggers), is a herbaceous plant that is indigenous to Asia, Europe, and North America. Since ancient times, this herb has been utilized for medicinal purposes. The therapeutic benefits of the plant are attributed to the phytochemicals found in its various sections.

A previous study found that *Taraxacum officinale* extracts have diuretic activity in a mouse model. And the ethanolic leaf extract of this plant increases urinary frequency and fluid excretion in healthy individuals. Additionally, two studies reported that compounds isolated from *Taraxacum officinale* exhibit hypoglycemic effects by inhibiting α -glucosidase and α -amylase.^[24]



Figure No:1.10 Taraxacum officinale

● URTICA DIOICA

Genus *Urtica*, or "nettle," is a genus of medicinal plants in the Urticaceae family that has been used medicinally since at least Ancient Greece. It has several health advantages.

Urtica dioica has traditionally been used as a diuretic in indigenous medicine. Experimentally, its aqueous extract exhibits natriuretic and diuretic activity in rabbits without affecting the rate of K^+ . Additionally, in a study on type 2 diabetic model rats, the water extract significantly lowered fasting serum glucose levels on the 14th day (14.05 ± 3.4 mmol/l in the control vs 8.3 ± 1.4 mmol/l in the treated group, $p = 0.04$). The extract also improved glycaemic and lipidemic status, likely

by affecting the histological and functional status of pancreatic β -cells.^[25,26]



Figure No:1.11 Urtica dioica

III. CONCLUSION

Diuretics are a cornerstone in the management of various cardiovascular and renal conditions. However, their use, especially with thiazide diuretics, often leads to the undesirable side effect of hyperglycemia. This poses a significant challenge in patients who are at risk of or managing diabetes mellitus, necessitating the exploration of alternative therapies that can mitigate these adverse effects.

The exploration of medicinal plants with dual diuretic and hypoglycemic activities offers a promising strategy for mitigating the hyperglycemia often induced by conventional diuretics. Plants such as *Samanea Saman*, *H. antidysentrica*, *Achyranthes Aspera*, *Taraxacum Officinale*, and *Urtica Dioica*, etc... have demonstrated significant potential in preclinical studies, showing efficacy in both increasing urine output and reducing blood glucose levels. The dual properties of these medicinal plants highlight their potential to create a more balanced therapeutic approach for patients requiring diuretic therapy but who are also at risk of hyperglycemia.

However, further clinical research is needed to validate these findings and ensure the safety and efficacy of these plants in human populations. This research will also help elucidate the mechanisms underlying their dual actions, paving the way for optimized treatment protocols.

Ultimately, the integration of these medicinal plants into conventional medicine could represent a significant advancement in the management of fluid retention and hyperglycemia, improving the quality of care for patients worldwide.

REFERENCES

- [1]. Wile D. Diuretics: a review. *Annals of clinical biochemistry*. 2012 Sep;49(5):419-31.
- [2]. Angappan R, Devanesan AA, Thilagar S.

- Diuretic effect of chlorogenic acid from traditional medicinal plant *Merremia emarginata* (Burm. F.) and its by-product hippuric acid. *Clinical Phytoscience*. 2018 Dec;4(1):1-6.
- [3]. Melka AE, Makonnen E, Debell A, et al. Diuretic activity of the aqueous crude extract and solvent fractions of the leaves of *Thymus serrulatus* in mice. *J Exp Pharmacol*. 2016;8:61-67.
 - [4]. Fathallah N, Slim R, Larif S, Hmouda H, Ben Salem C. Drug-induced hyperglycemia and diabetes. *Drug safety*. 2015 Dec; 38:1153-68.
 - [5]. Mandal AK, Hiebert LM. Is diuretic-induced hyperglycemia reversible and inconsequential? *Journal of Diabetes Research and Clinical Metabolism*. 2012 Jul 12;1(1):4.
 - [6]. Adverse reactions to bendroflumazide and propranolol for the treatment of mild hypertension. Report of Medical Research Council Working Party on Mild to Moderate Hypertension. *Lancet*. 1981; 2:539-543.
 - [7]. Elliott WJ, Meyer PM. Incident diabetes in clinical trials of antihypertensive drugs: a network meta-analysis. Meta-analysis of clinical trials showing that all antihypertensives analyzed, beta blockers, and thiazide diuretics are associated with the highest risk of diabetes. *Lancet* 2007;369(9557):201-7.
 - [8]. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA*. 2002; 288:2981-97.
 - [9]. Hirst JA, Farmer AJ, Feakins BG, Aronson JK, Stevens RJ. Quantifying the effects of diuretics and beta-blockers on glycaemic control in diabetes mellitus—a systematic review.
 - [10]. Ayvaz G, Balos T, Öner F, Karakoc A, Yetkin I, Cakir N, Arslan M. Acute and chronic effects of different concentrations of free fatty acids on the insulin-secreting function of islets. *Diabetes Metab*. 2002; 28:357-12.
 - [11]. Eriksson JW, Jansson PA, Carlberg B, Hagberg A, Kurland L, Svensson MK, et al. Hydrochlorothiazide, but not candesartan, aggravates insulin resistance and causes visceral and hepatic fat accumulation: the Mechanisms for the Diabetes Preventing Effect of Candesartan (MEDICA) study. *Hypertension*. 2008;52(6):1030-7.
 - [12]. Chukwuma CI, Matsabisa MG, Ibrahim MA, Erukainure OL, Chabalala MH, Islam MS. Medicinal plants with concomitant anti-diabetic and anti-hypertensive effects as potential sources of dual acting therapies against diabetes and hypertension: A review. *Journal of Ethnopharmacology*. 2019 May 10; 235:329-60.
 - [13]. Alam K, Hoq O, Uddin S. Medicinal plant *Allium sativum*. A review. *Journal of Medicinal Plant Studies*. 2016 Oct;4(6):72-9.
 - [14]. Patel MK, Rajput AP. Therapeutic significance of *Abutilon indicum*: An overview. *Am. J. Pharm. Tech. Res*. 2013; 4:20-35.
 - [15]. Al-Snafi AE. The pharmacological importance of *Benincasa hispida*. A review. *Int Journal of Pharma Sciences and Research*. 2013 Dec;4(12):165-70.
 - [16]. Mishra S, Aeri V, Gaur PK, Jachak SM. Phytochemical, therapeutic, and ethnopharmacological overview for a traditionally important herb: *Boerhavia diffusa* Linn. *BioMed research international*. 2014;2014(1):808302.
 - [17]. Prakash V, Jaiswal NI, Srivastava MR. A review on medicinal properties of *Centella asiatica*. *Asian J Pharm Clin Res*. 2017;10(10):69-74. and Al-Huda E, Debnath J. Evaluation of diuretic activity of aqueous extract of leaves of *Centella asiatica*.
 - [18]. Manvar M, Desai T. Phytochemical and pharmacological profile of *Ipomoea aquatica*. *Indian journal of medical sciences*. 2013 Mar 1;67(3/4):49.
 - [19]. Snigdha M, Kumar SS, Sharmistha M, Lalit S, Tanuja S. An overview on herbal medicines as diuretics with scientific evidence. *Scholars Journal of Applied Medical Science*. 2013 Oct 2;1(3):209-14.
 - [20]. Fan W, Fan L, Peng C, Zhang Q, Wang L, Li L, Wang J, Zhang D, Peng W, Wu C. Traditional uses, botany, phytochemistry, pharmacology, pharmacokinetics and toxicology of *Xanthium strumarium* L.: A

- review. *Molecules*. 2019 Jan 19;24(2):359.
- [22]. Vinodhini S. Review on ethnomedical uses, pharmacological activity and phytochemical constituents of *Samanea saman* (Jacq.) Merr. rain tree. *Pharmacognosy Journal*. 2018;10(2).
- [23]. Sinha S, Sharma A, Reddy PH, Rath B, Prasad NV, Vashishtha A. Evaluation of phytochemical and pharmacological aspects of *Holarrhena antidysenterica* (Wall.): A comprehensive review. *Journal of Pharmacy research*. 2013 Apr 1;6(4):488-92.
- [24]. Goyal BR, Goyal RK, Mehta AA. PHCOG rev.: plant review phyto- pharmacology of *Achyranthes aspera*: a review. *Pharmacogn Rev*. 2007 Jan;1(1):143-50.
- [25]. Di Napoli A, Zucchetti P. A comprehensive review of the benefits of *Taraxacum officinale* on human health. *Bulletin of the National Research Centre*. 2021 Jun 9;45(1):110.
- [26]. Das M, Sarma BP, Khan AK, Mosihuzzaman M, Nahar N, Ali L, Bhoulmik A, Rokeya B. The antidiabetic and antilipidemic activity of aqueous extract of *Urtica dioica* L. on type 2 diabetic model rats. *Journal of Bio- Science*. 2009; 17:1-6.
- [27]. Taheri Y, Quispe C, Herrera- Bravo J, Sharifi-Rad J, Ezzat SM, Merghany RM, Shaheen S, Azmi L, Prakash Mishra A, Sener B, Kılıç M. *Urtica dioica*-derived phytochemicals for pharmacological and therapeutic applications. *Evidence-based Complementary and Alternative Medicine*. 2022;2022(1):4024331.
- [28]. Shekshavali T, Roshan S. Evaluation for Diuretic Activity of *Abutilon indicum* and *Amaranthus spinosus* Leaves Extracts. *Res Rev a J Toxicol [Internet]*. 2017;7(2):12-5.
- [29]. Chude MA, Orisakwe OE, Afonne OJ, Gamaniel KS, Vongtau OH, Obi E. Hypoglycaemic effect of the aqueous extract of *Boerhavia diffusa* leaves. *Indian Journal of Pharmacology*. 2001 May 1;33(3):215-6.
- [30]. Manish Kumar Maity, Mamta Naagar, "Autoimmune Neurogenic Dysphagia", *International Journal of Science and Research (IJSR)*, Volume 11 Issue 7, July 2022, pp. 447-463, <https://www.ijsr.net/getabstract.php?paperid=SR22630151732>.
- [31]. Manish Kumar Maity, Mamta Naagar, "A Review on Headache: Epidemiology, Pathophysiology, Classifications, Diagnosis, Clinical Management and Treatment Modalities", *International Journal of Science and Research (IJSR)*, Volume 11 Issue 7, July 2022, pp. 506-515, <https://www.ijsr.net/getabstract.php?paperid=SR22703111804>.
- [32]. Md Shamshir Alam, Manish Kumar Maity, Abdul Salam Nazmi, Md Sarfaraz Alam, Md Salahuddin Ansari. Oral Health Issues And Preventive Measures In Geriatric Populations. *Journal of Pharmaceutical Negative Results [Internet]*. 2022 Dec. 31 [cited 2023 Jun. 24];:2647-55. Available from: <https://www.pnrjournal.com/index.php/home/article/view/9175>
- [33]. Nikita Sharma, Md Shamshir Alam, Anubha Sharma, Sanyam Garg, Manish Kumar Maity. Colorectal Cancer In Young Adults: Epidemiology, Risk Factors, Development, Symptoms, Traditional Herbal Therapy And Prevention. *Journal of Pharmaceutical Negative Results [Internet]*. 2022 Dec. 31 [cited 2023 Jun. 24];:1370-82. Available from: <https://pnjournal.com/index.php/home/article/view/6991>
- [34]. Ehteshamul Haque, Faiz Ahmed, Priyanka Chaurasiya, Neha Yadav, Nikita Dhiman, Manish Kumar Maity. (2023). A REVIEW ON ANTIDEPRESSANT EFFECT OF HERBAL DRUGS. *Journal of Pharmaceutical Negative Results*, 2716–2723. <https://doi.org/10.47750/pnr.2023.14.S02.319>.
- [35]. Omveer Singh, Shailesh Sharma, Mamta Naagar, Manish Kumar Maity, Eletriptan As Treatment Option For Acute Migraine, *International Journal Of Innovations & Research Analysis (Ijira)*, 02, 03(II), September, 2022, Pp 15-24.
- [36]. Priyanka Tanwar, Mamta Naagar, and Manish Kumar Maity, "Relationship between Type 2 Diabetes Mellitus and Osteoarthritis," *International Research Journal of Pharmacy and Medical Sciences (IRJPMS)*, Volume 6, Issue 2, pp. 59-70, 2023 (PDF) Relationship between Type 2 Diabetes Mellitus and Osteoarthritis. Available from: https://www.researchgate.net/publication/369022995_Relationship_between_Type_2_Diabetes_Mellitus_and_Osteoarthritis [accessed Jun 23 2023].
- [37]. Omveer Singh, Shailesh Sharma, Mamta Naagar, Manish Kumar Maity, Oral And Parenteral To Minimize The Nasal Delivery By Thermoreversible

- Mucoadhesive –A Review, International Journal Of Creative Research Thoughts (Ijcrt), 09/2022,10(9) Pp.-356-371.
- [38]. Md Shamshir Alam, Garima Malik, Priyanka Tanwar, Mamta Naagar, Tarun Singh, Omveer Singh, Manish Kumar Maity, A Review on Small-Cell Lung Cancer: Epidemiology, Pathophysiology, RiskFactors, Diagnosis, Clinical Management and Treatment Modalities, International Journal of Current Science Research and Review (ijcsrr), 06(01): 129-151.
- [39]. Priyanka Tanwar, Mamta Naagar, and Manish Kumar Maity, “Relationship between Diabetes Mellitus and Bone Health – A Review,”International Research Journal of Pharmacy and Medical Sciences (IRJPMS), Volume 6, Issue 2, pp. 46-58, 2023. (PDF) Relationship between Diabetes Mellitus and Bone Health - A Review. Available from: https://www.researchgate.net/publication/369022910_Relationship_between_Diabetes_Mellitus_and_Bone_Health_-_A_Review [accessed Jun 23 2023].
- [40]. Manish Kumar Maity. A review on Helicobacter pylori Infection. ijmsdr [Internet]. 2022Sep.17 [cited 2023Jun.23];6(9). Available from: <https://www.ijmsdr.com/index.php/ijmsdr/article/view/950>
- [41]. Md Shamshir Alam , Manish Kumar Maity , Abdul Salam Nazmi , Md Sarfaraz Alam , Md Salahuddin Ansari (2022) “Oral Health Issues And Preventive Measures In Geriatric Populations”,Journal of Pharmaceutical Negative Results, pp. 2647–2655. doi: 10.47750/pnr.2022.13.S10.316.